

Fluctuations of the Condensate in Ideal and Interacting Bose Gases

Hongwei Xiong^{1,2}, Shujuan Liu¹, Guoxiang Huang³, Zhijun Xu¹, Cunyuan Zhang¹

¹*Department of Applied Physics, Zhejiang University of Technology, Hangzhou, 310032, P. R. China*

²*Zhijiang College, Zhejiang University of Technology, Hangzhou, 310012, P. R. China*

³*Department of Physics, East China Normal University, 200062, Shanghai, P. R. China*

(February 1, 2008)

We investigate the fluctuations of the condensate in the ideal and weakly interacting Bose gases confined in a box of volume V within canonical ensemble. Canonical ensemble is developed to describe the behavior of the fluctuations when different methods of approximation to the weakly interacting Bose gases are used. Research shows that the fluctuations of the condensate exhibit anomalous behavior for the interacting Bose gas confined in a box.

03.75.Fi, 05.30.Jp

I. INTRODUCTION

The experimental achievement of Bose-Einstein condensation (BEC) in dilute alkali atoms [1], spin-polarized hydrogen [2] and recently in metastable helium [3] has enormously stimulated the theoretical research [4] on the ultracold bosons. In particular, fluctuations $\langle \delta^2 N_0 \rangle$ of the mean ground state occupation number N_0 have been recently thoroughly investigated in a series of papers. Apart from the intrinsic theoretical interest, it is foreseeable that such fluctuations will become experimentally testable in the near future [5].

It is well known that within grand canonical ensemble the fluctuations of the condensate are given by $\langle \delta^2 N_0 \rangle = N_0(N_0 + 1) \sim V^2$, implying that δN_0 becomes of order N when the temperature approaches zero. To avoid this sort of unphysically large condensate fluctuations, canonical (or microcanonical) ensemble has to be used to investigate the fluctuations of the condensate. Within microcanonical and canonical ensemble, the fluctuations of the condensate have been studied in a systematic way in the case of the ideal Bose gas [6–13]. Recently, the question of how interatomic interactions affect the fluctuations of the condensate has been the object of several theoretical investigations [14–19]. Giorgini *et al.* [14] found the anomalous behavior of the fluctuations in a weakly interacting Bose gas confined in a box within the traditional particle-number-nonconserving Bogoliubov approach. In [14] the fluctuations of the condensate follows the law $\langle \delta^2 N_0 \rangle \sim V^{4/3}$. However, Idziaszek *et al.* [15] considered that the fluctuations are proportional to the volume. Recently, Kocharovskiy *et al.* [18] supported and extended the results of the work of Giorgini *et al.* [14] using the particle-number-conserving operator formalism.

Although the correction to the ground state occupation number due to interatomic interaction has been clearly discussed within grand canonical ensemble [20] and canonical ensemble [21], the role of interaction on the condensate fluctuations of the weakly interacting Bose gas is still an open and unsolved problem. Different from the ground state occupation number, different models of describing the weakly interacting Bose gas will lead to vastly different prediction concerning the fluctuations of the condensate.

The purpose of this paper is to present a unified method to calculate the fluctuations of the condensate when different ways of approximation to the weakly interacting Bose gases are used. Within the canonical ensemble we give the distribution function of the ground state occupation number for the ideal and interacting Boson system in a box. We obtain the fluctuations of the condensate from the distribution function. In particular, we found that the distribution function is not Gaussian function in the case of interacting Boson system in a box. The paper is organized as follows. Sec.II is devoted to outline the canonical ensemble, which is developed to discuss the fluctuations of the condensate for the ideal Bose gas in a box. In Sec.III we investigate the fluctuations of the interacting Bose gas based on the lowest order perturbation theory. In Sec.IV the fluctuations are calculated based on the Bogoliubov theory. Finally, we give a discussion and summary of the results in Sec.V.

II. MEAN GROUND STATE OCCUPATION NUMBER AND FLUCTUATIONS IN THE IDEAL BOSE GASES

Let us start our investigation on the fluctuations of the ideal Bose gases in the frame of canonical ensemble. According to the canonical ensemble the partition function of the N non-interacting bosons in a box is given by

$$Z_{ideal}[N] = \sum_{\Sigma N_{\mathbf{n}}=N} \exp[-\beta \{\Sigma N_{\mathbf{n}} \varepsilon_{\mathbf{n}}\}], \quad (1)$$

where $N_{\mathbf{n}}$ and $\varepsilon_{\mathbf{n}}$ are the occupation numbers and energy level of the state $\mathbf{n} = \{n_x, n_y, n_z\}$ respectively. $\beta = 1/k_B T$. In (1) the energy level of the system takes the form

$$\varepsilon_{\mathbf{n}} = \frac{\pi^2 \hbar^2 (n_x^2 + n_y^2 + n_z^2)}{2mL^2}. \quad (2)$$

Separating out the ground state $\mathbf{n} = \mathbf{0}$ from the state $\mathbf{n} \neq \mathbf{0}$, we have

$$Z_{ideal}[N] = \sum_{N_0=0}^N \{ \exp[-\beta N_0 \varepsilon_0] Z_0(N - N_0) \}, \quad (3)$$

where $Z_0(N - N_0)$ stands for the partition function of a fictitious system comprising $N - N_0$ non-interacting bosons. Assuming $A_0(N - N_0)$ stands for the free energy of the fictitious system,

$$A_0(N - N_0) = -k_B T \ln Z_0(N - N_0) \quad (4)$$

From (3) and (4) the partition function $Z_{ideal}[N]$ becomes

$$Z_{ideal}[N] = \sum_{N_0=0}^N \exp[q(N, N_0)], \quad (5)$$

where $q(N, N_0) = -\beta N_0 \varepsilon_0 - \beta A_0(N - N_0)$. Obviously $\exp[q(N, N_0)]/Z_{ideal}[N]$ represents the probability to find N_0 atoms in the condensate. We will give the distribution function of the ground state occupation number in the following.

Let us first investigate the largest term in the sum of the partition function $Z_{ideal}[N]$. Assume the number of the condensed atoms is N_0^p in the largest term of the partition function $Z_{ideal}[N]$. The largest term $Z_0(N - N_0^p)$ is determined by the requirement that $\frac{\partial}{\partial N_0} q(N, N_0)|_{N_0=N_0^p} = 0$, *ie.*

$$-\beta \varepsilon_0 - \beta \frac{\partial}{\partial N_0^p} A_0(N - N_0^p) = 0 \quad (6)$$

The calculations of the free energy $A_0(N - N_0^p)$ is non-trivial because there is a requirement that the number of the particles is $N - N_0^p$ in the summation of the partition function $Z_0(N - N_0^p)$. Using the saddle-point method developed by Darwin and Fowler [22] it is straightforward to obtain the free energy $A_0(N - N_0^p)$ of the fictitious system.

$$A_0(N - N_0^p) = (N - N_0^p) k_B T \ln z_0^p - V \frac{k_B T}{\lambda^3} g_{5/2}(z_0^p), \quad (7)$$

where $\lambda = \sqrt{2\pi\beta\hbar^2/m}$ is the thermal wavelength. z_0^p is the fugacity of the $N - N_0^p$ non-interacting bosons and is determined by the equation

$$N - N_0^p = \sum_{\mathbf{n} \neq \mathbf{0}} \frac{1}{\exp[\varepsilon_{\mathbf{n}}/k_B T] (z_0^p)^{-1} - 1} = \frac{V}{\lambda^3} g_{3/2}(z_0^p). \quad (8)$$

From (7) and (8) one finds

$$-\beta \frac{\partial}{\partial N_0^p} A_0(N - N_0^p) = \ln z_0^p. \quad (9)$$

Combining (6) and (9) one obtains $\ln z_0^p = \beta \varepsilon_0$. Therefore the most probable value N_0^p is determined by

$$N_0^p = N - \sum_{\mathbf{n} \neq \mathbf{0}} \frac{1}{\exp[(\varepsilon_{\mathbf{n}} - \varepsilon_0)/k_B T] - 1}. \quad (10)$$

N_0^p is exactly the mean occupation number of the condensate atoms in the frame of the grand canonical ensemble. For sufficiently large N , the sum $\sum_{N_0=0}^N$ in (3) maybe replaced by the largest term, for the error omitted in doing so will be statistically negligible. In this case, (10) shows the equivalence between canonical ensemble and grand canonical ensemble for large N . From (10), below the critical temperature N_0^p is determined by

$$N_0^p = N - \frac{V}{\lambda^3} \zeta(3/2) = N \left(1 - \left(\frac{T}{T_c^0} \right)^{3/2} \right), \quad (11)$$

where $T_c^0 = \frac{2\pi}{[\zeta(3/2)]^{2/3}} \frac{\hbar^2}{mk_B} \left(\frac{N}{V} \right)^{2/3}$ is the transition temperature of the ideal Bose gas.

Other terms in (5) will contribute to the fluctuations of the condensate, and lead to the deviation of the mean occupation number $\langle N_0 \rangle$ from the most probable value N_0^p . When $N_0 \neq N_0^p$, $\frac{\partial}{\partial N_0} q(N, N_0) \neq 0$. Assuming

$$\frac{\partial}{\partial N_0} q(N, N_0) = \alpha(N, N_0), \quad (12)$$

and repeating the saddle-point method, one obtains

$$N_0 = N - \sum_{\mathbf{n} \neq \mathbf{0}} \frac{1}{\exp[(\varepsilon_{\mathbf{n}} - \varepsilon_0)/k_B T] \exp[-\alpha(N, N_0)] - 1}. \quad (13)$$

From (11) and (13) one gets

$$\alpha(N, N_0) = -\frac{\lambda^6}{V^2} \frac{(N_0 - N_0^p)^2}{4\pi} \theta(N_0 - N_0^p) \quad (14)$$

where $\theta(N_0 - N_0^p)$ is a Sign function. $\theta(N_0 - N_0^p) = 1$ when $N_0 > N_0^p$, and $\theta(N_0 - N_0^p) = -1$ when $N_0 < N_0^p$. To obtain (14) we have used the expansions $g_{3/2}(1 - \delta) \approx \zeta(3/2) - 2\sqrt{\pi}\delta$ [23] and the approximation $\exp[-\alpha(N, N_0)] \approx 1 - \alpha(N, N_0)$. From (12) and (14) one obtains easily the following result for $q(N, N_0)$.

$$\begin{aligned} q(N, N_0) &= \int_{N_0^p}^{N_0} \alpha(N, N_0) dN_0 + q(N, N_0^p) \\ &= -\frac{\lambda^6}{12\pi V^2} |N_0 - N_0^p|^3 + q(N, N_0^p). \end{aligned} \quad (15)$$

The partition function $Z_{ideal}[N]$ is thus

The coefficient in (21) is $A = 2/\pi^4 \times \sum_{\mathbf{n} \neq \mathbf{0}} 1/\mathbf{n}^4 = 0.105$. The dashed line is larger than our result because of the approximation in [14]. In [14] $\langle \delta^2 N_{\mathbf{0}} \rangle = \sum_{\mathbf{n} \neq \mathbf{0}} f_{\mathbf{n}}^2$, where $f_{\mathbf{n}} = (\exp(\varepsilon_{\mathbf{n}}/k_B T) - 1)^{-1}$. For the convenience of calculations, $f_{\mathbf{n}}$ is approximated as $\varepsilon_{\mathbf{n}}/k_B T$ for low energy atoms. However, this approximation is used also for the atoms whose energy level is larger than $k_B T$. Obviously, this approximation will lead to the fluctuations become larger. On the other hand, (21) holds in the canonical ensemble except near and above T_c , while our analysis holds also for the temperature near T_c^0 . In Fig.2 the dotted line shows the numerical result of Wilkens *et al.* [9].

In Fig.2 the arrow marks the temperature T_m which corresponds to the maximum fluctuations $\langle \delta^2 N_{\mathbf{0}} \rangle_{\max}$. Below the temperature T_m , from (19), one obtains the analytical result for the fluctuations of the condensate.

$$\langle \delta^2 N_{\mathbf{0}} \rangle = A \left(\frac{mk_B T}{\hbar^2} \right)^2 V^{4/3}. \quad (22)$$

It is interested to find that the coefficient differs by a factor 2, compared to (21). The thin solid line shows (22) in Fig.2.

We should note that our results are reliable although the disputable saddle-point method is used to investigate the fluctuations of the condensate. It is well known that the applicability of the saddle-point approximation for the condensed Bose gases has been the subject of a long debate [7,24]. Recently, the analysis in [12] showed that the fluctuations are overestimated, and do not appear to vanish properly with temperature using the usual saddle-point method. Our discussions for fluctuations are reasonable because of two reasons. (i) As proved in [13], the free energy (7) of the non-interacting Bose gases is still correct, even when carefully deal with the failure of the standard saddle-point method below the critical temperature. (ii) In the usual statistical method $\langle N_{\mathbf{0}} \rangle$ and $\langle \delta^2 N_{\mathbf{0}} \rangle$ are obtained through the first and second partial derivative of the partition function respectively. When saddle-point approximation is used to calculate the partition function of the system, the error will be overestimated in the second partial derivative of the partition function so that we can not obtain correct fluctuations of the condensate in the usual method. However, in this paper what we used is the reliable result (7). The distribution function of the ground state occupation number is obtained directly from (7), without resorting to the second partial derivative of the partition function. $\langle N_{\mathbf{0}} \rangle$ and $\langle \delta^2 N_{\mathbf{0}} \rangle$ are obtained from the distribution function in this paper.

III. FLUCTUATIONS OF THE CONDENSATE BASED ON THE LOWEST ORDER PERTURBATION THEORY

In the case of interacting Bose gases, the role of interactions on the fluctuations of the condensate is still an open and unsolved problem. Giorgini *et al.* [14] predicted the anomalous behavior of the fluctuations in a weakly interacting Bose gas confined in a box, while Idziaszek *et al.* [15] considered that the fluctuations are normal. Researches show that different model of approximation to the interacting Bose gases will lead to different predictions concerning the fluctuations of the condensate. The method developed to obtain the fluctuations of the ideal Bose gas in this paper can be used straightforwardly to discuss the fluctuations of the interacting Bose gases when different models of approximation are adopted.

Let us first discuss the fluctuations of the condensate in the case of the lowest order perturbation theory, which is also discussed in [15]. In terms of the lowest order perturbation theory, the partition function of the system within the canonical ensemble is given by

$$Z_{int}[N] = \sum_{\Sigma N_{\mathbf{n}}=N} \exp[-\beta(\Sigma N_{\mathbf{n}} \varepsilon_{\mathbf{n}} + E_{int})], \quad (23)$$

where the interaction energy of the system takes the form [25,26]

$$E_{int} = \frac{4\pi a \hbar^2}{mV} \left(N^2 - \frac{1}{2} N_{\mathbf{0}}^2 \right). \quad (24)$$

In (24) a is the scattering length. Separating out the ground state $\mathbf{n} = \mathbf{0}$ from the state $\mathbf{n} \neq \mathbf{0}$, one obtains the following form for the partition function

$$Z_{int}[N] = \sum_{N_{\mathbf{0}}=0}^N \{ \exp[-\beta N_{\mathbf{0}} \varepsilon_{\mathbf{0}} - \beta E_{int}] Z_0(N - N_{\mathbf{0}}) \}, \quad (25)$$

where $Z_0(N - N_{\mathbf{0}})$ stands for the partition function of a fictitious $N - N_{\mathbf{0}}$ non-interacting bosons. Using the free energy $A_0(N - N_{\mathbf{0}})$ of the fictitious system, the partition function is thus

$$Z_{int}[N] = \sum_{N_{\mathbf{0}}=0}^N \exp[q(N, N_{\mathbf{0}})], \quad (26)$$

where $q(N, N_{\mathbf{0}})$ takes the form

$$q(N, N_{\mathbf{0}}) = -\beta N_{\mathbf{0}} \varepsilon_{\mathbf{0}} - \beta E_{int} - \beta A_0(N - N_{\mathbf{0}}). \quad (27)$$

Analogously to the case of ideal Bose gases, let us first investigate the largest term in the sum of $Z_{int}[N]$. The largest term is determined by the requirement $\frac{\partial}{\partial N_{\mathbf{0}}} q(N, N_{\mathbf{0}})|_{N_{\mathbf{0}}=N_{\mathbf{0}}^p} = 0$. Therefore one obtains the most probable value $N_{\mathbf{0}}^p$ of the interacting Bosons.

easy to recover the fluctuations of the ideal Bose gases. According to (39) the leading contributions to the fluctuations are anomalous, *ie.* proportional to $N^{4/3}$, while there are also normal contributions proportional to N due to interatomic interaction. This conclusion contradicts with that of [15] which predicts normal behavior, where the lowest order perturbation theory is also used to discuss the fluctuations of the condensate.

IV. FLUCTUATIONS OF THE CONDENSATE BASED ON BOGOLIUBOV THEORY

Let us investigate the fluctuations of the condensate in the framework of Bogoliubov theory of a uniform weakly interacting Bose gas confined in a box. According to Bogoliubov theory [27,28], the total number of particles out of the condensate is given by

$$N_T = \sum_{\mathbf{n} \neq 0} N_{\mathbf{n}} = \sum_{\mathbf{n} \neq 0} (u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2) f_{\mathbf{n}}, \quad (40)$$

where

$$u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2 = \frac{\left((\varepsilon_{\mathbf{n}}^B)^2 + g^2 n_0^2\right)^{1/2}}{2\varepsilon_{\mathbf{n}}^B}, \quad (41)$$

$$u_{\mathbf{n}} v_{\mathbf{n}} = -\frac{g n_0}{2\varepsilon_{\mathbf{n}}^B}, \quad (42)$$

and $f_{\mathbf{n}}$ is the number of quasi-particles present in the system at the thermal equilibrium.

$$f_{\mathbf{n}} = \frac{1}{\exp[\varepsilon_{\mathbf{n}}^B/k_B T] - 1} \quad (43)$$

In addition, the energy of the quasi-particles entering (41) and (42) is given by the well known Bogoliubov spectrum

$$\varepsilon_{\mathbf{n}}^B = \left((\varepsilon_{\mathbf{n}} + g n_0)^2 - g^2 n_0^2\right)^{1/2}, \quad (44)$$

where $g = 4\pi\hbar^2 a/m$ is the coupling constant, and $n_0 = N_0/V$ is the condensate density. At low $|\mathbf{n}| = \sqrt{n_x^2 + n_y^2 + n_z^2}$, one obtains $u_{\mathbf{n}}^2 \simeq v_{\mathbf{n}}^2 \propto 1/|\mathbf{n}|$ and $f_{\mathbf{n}} \propto 1/|\mathbf{n}|$. This results in $1/|\mathbf{n}|^2$ divergence in (43) at low $|\mathbf{n}|$. Although this sort of divergence will not lead to large contribution to the number of low energy quasi-particles, it gives leading contribution to the fluctuations of the condensate, as pointed out in [14]. We will investigate the fluctuations due to low energy quasi-particles in the following.

In (40) $N_{\mathbf{n}}$ can be regarded as the effective occupation number of the thermal atoms, while

$$N_{\mathbf{n}}^B = \frac{N_{\mathbf{n}}}{u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2} = f_{\mathbf{n}} \quad (45)$$

is the occupation number of the quasi-particles. From the form of $f_{\mathbf{n}}$, we can construct the partition function of the quasi-particles in the frame of canonical ensemble.

$$Z_B = \sum_{\{\mathbf{n}\}} \exp \left[-\beta \sum_{\mathbf{n}} N_{\mathbf{n}}^B \varepsilon_{\mathbf{n}}^B \right]. \quad (46)$$

From (45) Z_B becomes

$$Z_B = \sum_{\{\sum N_{\mathbf{n}} = N\}} \exp \left[-\beta \sum_{\mathbf{n}} N_{\mathbf{n}} \varepsilon_{\mathbf{n}}^{eff} \right], \quad (47)$$

where $\varepsilon_{\mathbf{n}}^{eff} = \varepsilon_{\mathbf{n}}^B / (u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2)$ can be regarded as an effective energy level of the thermal atoms. In this case Z_B is the partition function of a fictitious Boson system comprising N non-interacting Bosons whose energy level is determined by $\varepsilon_{\mathbf{n}}^{eff}$. From (47) the most probable value N_0^p is given by

$$N_0^p = N - \sum_{\mathbf{n} \neq 0} \frac{1}{\exp \left[(\varepsilon_{\mathbf{n}}^{eff} - \varepsilon_0^{eff}) / k_B T \right] - 1}, \quad (48)$$

Obviously the occupation number of low $|\mathbf{n}|$ in (48) coincides with that of (40). Analogously, other N_0 is thus

$$N_0 = N - \sum_{\mathbf{n} \neq 0} \frac{1}{\exp \left[(\varepsilon_{\mathbf{n}}^{eff} - \varepsilon_0^{eff}) / k_B T \right] \exp[-\alpha(N, N_0)] - 1}. \quad (49)$$

From (48) and (49) one gets

$$\alpha(N, N_0) \approx -\frac{N_0 - N_0^p}{\sum_{\mathbf{n} \neq 0} (u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2)^2 f_{\mathbf{n}}^2}, \quad (50)$$

where we have used the approximation $f_{\mathbf{n}} \approx k_B T / \varepsilon_{\mathbf{n}}^B$ for low energy quasi-particles. Therefore the Gaussian distribution function of the system is given by

$$G_B(N, N_0) = \exp \left[-\frac{(N_0 - N_0^p)^2}{2 \sum_{\mathbf{n} \neq 0} (u_{\mathbf{n}}^2 + v_{\mathbf{n}}^2)^2 f_{\mathbf{n}}^2} \right] \\ \approx \exp \left[-\frac{(\zeta(3/2))^{4/3} (N_0 - N_0^p)^2}{(2\pi)^2 A N^{4/3} t^2} \right]. \quad (51)$$

Obviously the mean occupation number $\langle N_0 \rangle$ and fluctuations $\langle \delta^2 N_0 \rangle$ is given by

$$\langle N_0 \rangle = \frac{\sum_{N_0=0}^N N_0 G_B(N, N_0)}{\sum_{N_0=0}^N G_B(N, N_0)}, \quad (52)$$

- S C, Kleppner D and Greytak T J 1998 *Phys. Rev. Lett.* **81** 3811
- [3] Robert A, Sirjean O, Browaeys A, Poupard J, Nowak S, Boiron D, Westbrook C I, Aspect A 2001 *Science* **292** 461
Pereira Dos Santos F, Leonard J, Wang J, Barrelet C J, Perales F, Rasel E, Unnikrishnan C S, Leduc M, Cohen-Tannoudji C 2001 *Phys. Rev. Lett.* **86** 3459
- [4] Dalfovo F, Giorgini S, Pitaevskii L P and Stringari S 1999 *Rev. Mod. Phys.* **71** 463
- [5] Idziaszek Z, Rzażewski K and Lewenstein M 2000 *Phys. Rev. A* **61** 053608
- [6] Hauge E H 1969 *Physica Norvegica* **4** 19
Fujiwara I, ter Haar D and Wergeland H 1970 *J. Stat. Phys.* **2** 329
- [7] Ziff R M, Uhlenbeck G E and Kac M 1977 *Phys. Rep.* **32** 169
- [8] Borrmann P, Harting J, Mülken O and Hilf E R 1999 *Phys. Rev. A* **60**, 1519
- [9] Wilkens M and Weiss C 1997 *J. Mod. Opt.* **44** 1801
- [10] Politzer H D 1996 *Phys. Rev. A* **54** 5048
- [11] Gajda M and Rzażewski K 1997 *Phys. Rev. Lett.* **78** 2686
Navez P, Bitouk D, Gajda M, Idziaszek Z and Rzażewski K 1997 *Phys. Rev. Lett.* **79** 1789
Holthaus M, Kalinowski E and Kirsten K 1998 *Ann. Phys. (N. Y.)* **270** 198
- [12] Grossmann S, Holthaus M 1997 *Phys. Rev. Lett.* **79** 3557
- [13] Holthaus M, Kalinowski E 1999 *Ann. Phys. (N. Y.)* **276** 321.
- [14] Giorgini S, Pitaevskii L P and Stringari S 1998 *Phys. Rev. Lett.* **80** 5040
- [15] Idziaszek Z, Gajda M, Navez P, Wilkens M and Rzażewski K 1999 *Phys. Rev. Lett.* **82** 4376
- [16] Illuminati F, Navez P and Wilkens M 1999 *J. Phys. B* **32** L461
- [17] Meier F and Zwerger W 1999 *Phys. Rev. A* **60** 5133
- [18] Kocharovsky V V, Kocharovsky V I V and Scully M O 2000 *Phys. Rev. Lett.* **84** 2306
- [19] Graham R 2000 *Phys. Rev. A* **62** 023609
- [20] Giorgini S, Pitaevskii L P and Stringari S 1996 *Phys. Rev. A* **54** R4633
Naraschewski M, Stamper-Kurn D M 1998 *Phys. Rev. A* **58** 2423
Liu S J, Huang G X, Ma L, Zhu S H and Xiong H W 2000 *J. Phys. B* **33** 3911
- [21] Xiong H W, Liu S J, Huang G X, Xu Z J and Zhang C Y 2001 *J. Phys. B* **34** 3013
- [22] Darwin C G and Fowler R H 1922 *Phil. Mag.* **44** 450, 823; 1922 *Proc. Cambridge Phil. Soc.* **21** 262
- [23] Robinson J E 1951 *Phys. Rev.* **83** 678
London F 1954 *Superfluids* (Wiley, New York) Vol. II Appendix p.203
- [24] Dingle R B 1949 *Proc. Cambridge. Phil. Soc* **45** 275
Fraser A R 1951 *Phil. Mag.* **42** 165
- [25] Pathria R K 1972 *Statistical Mechanics* (Pergamon Press: New York)
- [26] Huang K 1987 *Statistical Mechanics* (John Wiley and Sons, New York)
- [27] Bogoliubov N 1947 *J. Phys. USSR* **11** 23
- [28] Giorgini S, Pitaevskii L P and Stringari S 1997 *J. Low. Temp. Phys.* **109** 309